

For The Registered Medical Practitioner or a Hospital Use Only

Rx

Varenicline Tartrate Tablets 0.5 mg / 1 mg **VARENISMART 0.5 mg / 1 mg**

Composition:

Each Film Coated Tablet Contains:
Varenicline Tartrate 0.5 mg./1 mg
Excipients Q.S.
Colour: Titanium Dioxide BP

Indication

VARENISMART 0.5 MG / 1 MG contains the active substance varenicline. VARENISMART 0.5 MG / 1 MG is a medicine which is used in adults to help them stop smoking.
VARENISMART 0.5 MG / 1 MG can help to relieve the craving and withdrawal symptoms associated with stopping smoking.
VARENISMART 0.5 MG / 1 MG can also reduce the enjoyment of cigarettes if you do smoke when on treatment.

Warnings and precautions

Talk to your doctor or pharmacist before taking VARENISMART 0.5 MG / 1 MG
There have been reports of depression, suicidal ideation and behaviour and suicide attempts in patients taking VARENISMART 0.5 MG / 1 MG If you are taking VARENISMART 0.5 MG / 1 MG and develop agitation, depressed mood, changes in behaviour that are of concern to you or your family or if you develop suicidal thoughts or behaviours you should stop taking VARENISMART 0.5 MG / 1 MG and contact your doctor immediately for treatment assessment.

Children and adolescents

VARENISMART 0.5 MG / 1 MG is not recommended for use in paediatric patients as efficacy was not demonstrated.

Other medicines and varenismart

In some cases, as a result of stopping smoking, with or without VARENISMART 0.5 MG / 1 MG, an adjustment of the dose of other medicines may be necessary. Examples include theophylline (a medicine to treat breathing problems), warfarin (a medicine to reduce blood clotting), and insulin (a medicine to treat diabetes). If in doubt, you should consult your doctor or pharmacist.

If you have severe kidney disease you should avoid taking cimetidine (a medicine used for gastric problems) at the same time as VARENISMART 0.5 MG / 1 MG as this may cause increased blood levels of VARENISMART 0.5 MG / 1 MG

Use of VARENISMART 0.5 MG / 1 MG with other therapies for smoking cessation

Consult your doctor before using VARENISMART 0.5 MG / 1 MG in combination with other smoking cessation therapies.

VARENISMART 0.5 MG / 1 MG with food and drink

There have been some reports of increased intoxicating effects of alcohol in patients taking VARENISMART 0.5 MG / 1 MG However, it is not known if VARENISMART 0.5 MG / 1 MG actually increases alcohol intoxication.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine

Driving and using machines

VARENISMART 0.5 MG / 1 MG may be linked with dizziness, sleepiness and transient loss of consciousness. You should not drive, operate complex machinery or engage in any other potentially hazardous activities until you know whether this medicine affects your ability to perform these activities.

How to take

As direction by the physician.

Side effects

Chest infection, inflammation of the sinuses

Increased weight, decreased appetite, increased appetite

Sleepiness, dizziness, changes in the way things taste

Shortness of breath, cough

Heartburn, vomiting, constipation, diarrhoea, feeling bloated, abdominal pain, toothache, indigestion, flatulence, dry mouth

Storage:

Storage : Store in a cool & dry place,
at a temperature not exceeding 30°C.
Protected from light & moisture.

Packing : 10 x 1 x 10 Tablets

Marketed by :



Healing Pharma India Pvt. Ltd.
(An ISO 9001-2015 Certified Company)

B-411, Western Edge II Premises CHS Ltd,
Near Metro Mall, Western Express Highway,
Borivali (East). Mumbai - 400066

International Correspondence office

Healing Pharma LLC, DE USA 

"ACCORDING TO EU GMP STANDARD"