

R_x Sumatriptan Tablets BP

SUMITOP[®]

सुमीटोप

Composition:

Each film coated tablet contains:

Sumatriptan Succinate IP

Eq. to Sumatriptan25mg

Excipientsq.s.

Colour: Red oxide of iron & Titanium dioxide IP

Each film coated tablet contains:

Sumatriptan Succinate IP

Eq. to Sumatriptan50mg

Excipientsq.s.

Colour: Red oxide of iron & Titanium dioxide IP

Each film coated tablet contains:

Sumatriptan Succinate IP

Eq. to Sumatriptan100mg

Excipientsq.s.

Colour: Red oxide of iron & Titanium dioxide IP

Indications: Treatment of acute migraine; cluster headache

Cautions: 5HT₁-receptor agonists should be used with caution in the elderly [unlicensed], and in conditions which predispose to coronary artery disease (preexisting cardiac disease), history of seizures; sensitivity to sulfonamides.

Driving drowsiness may affect performance of skilled tasks (e.g. driving).

Contra-indications: 5HT₁-receptor agonists are contra-indicated in ischaemic heart disease, previous myocardial infarction, coronary vasospasm (including Prinzmetal's angina), and uncontrolled or severe hypertension. 5HT₁-receptor agonists are not indicated for the treatment of hemiplegic, basilar, or ophthalmoplegic migraine; previous cerebrovascular accident or transient ischaemic attack; peripheral vascular disease; moderate and severe hypertension.

Hepatic impairment: Reduce oral dose to 25–50 mg; avoid in severe impairment

Renal impairment: Use with caution.

Pregnancy: There is limited experience of using 5HT₁ receptor agonists during pregnancy; manufacturers advise that they should be avoided unless the potential benefit outweighs the risk.

Breast-feeding: Present in milk but amount probably too small to be

harmful; withhold breast-feeding for 12 hours.

Side-effects: Sensations of tingling, heat, heaviness, pressure, or tightness of any part of the body (including throat and chest—discontinue if intense, may be due to coronary vasoconstriction or to anaphylaxis), flushing, dizziness, feeling of weakness; fatigue; nausea and vomiting also reported; also dyspnoea, drowsiness, transient increase in blood pressure, myalgia; also reported diarrhoea, ischaemic colitis, hypotension, bradycardia or tachycardia, palpitation, arrhythmias, myocardial infarction, Raynaud's syndrome, anxiety, seizures, tremor, dystonia, nystagmus, arthralgia, visual disturbances, and sweating; epistaxis with nasal spray.

Dosage:

By mouth, migraine, 50mg (some patients may require 100 mg); dose may be repeated after at least 2 hours if migraine recurs; max. 300mg in a day.

Storage:

Store at a temperature not exceeding 30°C.

Protect from light.

Keep medicine out of reach of children.

Manufactured By:



Healing Pharma India Pvt. Ltd.

(An ISO 9001-2015 Certified Company)

Shop No. 01, Plot No.: 25-B,
Dev Industrial Estate, Gorwa, Vadodara-16.

Reg. Office: B-411,
Western Edge II Premises CHS Ltd,
Near Metro Mall, Western Express Highway,
Borivali (East). Mumbai-400066
International Correspondence office
Healing Pharma LLC, DE USA

Email : info@healingpharma.in

Customer Care No. : 7466030303

At. G/5 & G/6, Industrial Estate,
Gorwa, Vadodara-390016.